

* The temporal Heisenberg inequality

• Ehrenfest theorem:

$$\frac{d}{dt} \langle A \rangle_\psi = \frac{1}{i\hbar} \langle [A, H] \rangle_\psi \quad \parallel \langle \cdot \rangle_\psi = \langle \psi(t) | \cdot | \psi(t) \rangle$$

• uncertainty relation: $\langle (\Delta A)^2 \rangle_\psi \langle (\Delta B)^2 \rangle_\psi \geq \frac{1}{4} |\langle [A, B] \rangle_\psi|^2$

Let's put H into B ! $\equiv (\Delta H)^2$

$$\Rightarrow \Delta_\psi H \Delta_\psi A \geq \frac{1}{2} |\langle [A, H] \rangle_\psi| = \frac{1}{2} \hbar \left| \frac{d}{dt} \langle A \rangle_\psi \right|$$

If we define the time $\tau_\psi(A)$ as

$$\frac{1}{\tau_\psi(A)} \equiv \left| \frac{d\langle A \rangle_\psi}{dt} \right| \frac{1}{\Delta_\psi A},$$

then τ_ψ = characteristic time for ^{the} expectation value of A to change by $\Delta_\psi A$.

$$\Rightarrow \Delta_\psi H \tau_\psi(A) \geq \frac{1}{2} \hbar \Rightarrow \underbrace{\Delta E}_{\text{Energy spread}} \underbrace{\tau_\psi(A)}_{\text{characteristic evolution time}} \geq \frac{1}{2} \hbar$$

2.3 Simple Harmonic oscillator

(a) Energy eigenkets. (b) Dirac's operator method)

$$H = \frac{\tilde{p}^2}{2m} + \frac{1}{2} m \omega^2 \tilde{x}^2 = \hbar \omega \left(\tilde{a}^\dagger \tilde{a} + \frac{1}{2} \right)$$

$$\equiv \hbar \omega \left(\tilde{N} + \frac{1}{2} \right)$$

def. $\tilde{a} = \sqrt{\frac{m\omega}{2\hbar}} \left(\tilde{x} + i \frac{\tilde{p}}{m\omega} \right)$ $\Rightarrow \begin{cases} \tilde{x} = \frac{x_0}{\sqrt{2}} (\tilde{a} + \tilde{a}^\dagger) \\ \tilde{p} = i \frac{\hbar}{\sqrt{2} x_0} (-\tilde{a} + \tilde{a}^\dagger) \end{cases}$

creation operator $\tilde{a}^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(\tilde{x} - i \frac{\tilde{p}}{m\omega} \right)$

$\tilde{N} = \tilde{a}^\dagger \tilde{a}$

$\parallel x_0 = \sqrt{\frac{\hbar}{m\omega}}$

\Rightarrow Commutation relation $[\tilde{a}, \tilde{a}^\dagger] = 1$

$\Rightarrow [H, \tilde{N}] = 0$; There's simultaneous eigenkets

$$\therefore H|n\rangle = (n + \frac{1}{2})\hbar\omega |n\rangle \quad || \quad \tilde{N}|n\rangle = n|n\rangle$$

\nearrow same eigenkets $\{|n\rangle\}$.

$$\Rightarrow \boxed{E_n = (n + \frac{1}{2})\hbar\omega} \quad || \quad H|n\rangle = E_n|n\rangle$$

Q. What's "n", then? We know that $n = 0, 1, 2, \dots$

But, how do we know that
from what we have?

Let's look at the commutation relation:

$$\begin{aligned} \underline{[\tilde{N}, \tilde{a}]} &= [\tilde{a}^\dagger \tilde{a}, \tilde{a}] = \underbrace{\tilde{a}^\dagger [\tilde{a}, \tilde{a}]}_0 + \underbrace{[\tilde{a}^\dagger, \tilde{a}]}_{-1} \tilde{a} \\ &= \underline{-\tilde{a}} \end{aligned}$$

likewise, $\underline{[\tilde{N}, \tilde{a}^\dagger] = \tilde{a}^\dagger}$.

Now, try $\underline{\tilde{N} \tilde{a}^\dagger |n\rangle} = ([\tilde{N}, \tilde{a}^\dagger] + \tilde{a}^\dagger \tilde{N}) |n\rangle$
 $\quad \quad \quad \swarrow$
 $\quad \quad \quad \text{switch!}$
 $\quad \quad \quad = \tilde{a}^\dagger (\tilde{N} + 1) |n\rangle$

$$\Rightarrow \underline{\tilde{N} \tilde{a}^\dagger |n\rangle = (n+1) \tilde{a}^\dagger |n\rangle}$$

Thus, \tilde{a}^\dagger is a creation operator
as it makes $n \rightarrow n+1$

likewise, $\underline{\tilde{N} \tilde{a} |n\rangle} = ([\tilde{N}, \tilde{a}] + \tilde{a} \tilde{N}) |n\rangle$
 $\quad \quad \quad = \underline{(n-1) |n\rangle}$

$\Rightarrow \tilde{a}$ is an annihilation operator,
($n \rightarrow n-1$)

$$\left[\begin{array}{l} \tilde{N} \tilde{a}^+ |n\rangle = (n+1) \tilde{a}^+ |n\rangle \\ \tilde{N} \tilde{a} |n\rangle = (n-1) \tilde{a} |n\rangle \end{array} \right. \xrightarrow{\text{implies}}$$

$$\tilde{a}^+ |n\rangle = c_+ |n+1\rangle$$

$$\tilde{a} |n\rangle = c_- |n-1\rangle$$

* let's check: $\tilde{N} \tilde{a}^+ |n\rangle = (n+1) \tilde{a}^+ |n\rangle$ || c_{\pm} : c-number.

$$\Rightarrow \tilde{N} \cdot c_+ |n+1\rangle = (n+1) \cdot c_+ |n+1\rangle$$

$$\tilde{N} |n+1\rangle = (n+1) |n+1\rangle \quad : \quad 0 \leq n$$

Now, let's determine c_{\pm} .

To recover $\tilde{N} |n\rangle = n |n\rangle$,

$$\langle n | \tilde{a}^+ \tilde{a} |n\rangle = |c_-|^2 \langle n-1 | n-1\rangle = n$$

$$\therefore \underline{c_- = \sqrt{n}}$$

|| choose c_{\pm} to be real and positive.

$$\text{likewise, } \langle n | \tilde{a}^+ \tilde{a} |n\rangle = \langle n | \cdot \left(\tilde{a} \tilde{a}^+ - \underbrace{\tilde{a} \tilde{a}^+ + \tilde{a}^+ \tilde{a}}_{= [\tilde{a}^+, \tilde{a}] = -1} \right) \cdot |n\rangle$$

$$= \langle n | \tilde{a} \tilde{a}^+ |n\rangle - 1$$

$$\Rightarrow |c_+|^2 = n+1 \quad \therefore \underline{c_+ = \sqrt{n+1}}$$

Now, we know that, if $(n, |n\rangle)$ is the eigenpair,

$$\text{So ARE: } \{ \dots (n-2, |n-2\rangle), (n-1, |n-1\rangle), \\ (n+1, |n+1\rangle), (n+2, |n+2\rangle), \dots \}$$

Not enough to determine n .

Is there any lower bound?

$$\bullet \quad n = 0, 1, 2, 3, 4, \dots$$


$$\Rightarrow \langle n | \tilde{N} |n\rangle = \langle n | \tilde{a}^+ \rangle \cdot \langle a | n \rangle \geq 0$$

$$\therefore \underline{n \geq 0}$$

⇒ Ground-state energy

Eigen state

$$E_0 = \frac{1}{2} \hbar \omega$$



$$\begin{aligned} &|0\rangle \\ &\hookrightarrow |1\rangle \\ &\hookrightarrow |2\rangle \\ &\vdots \end{aligned}$$

$$|n+1\rangle = \frac{\tilde{a}^\dagger}{\sqrt{n+1}} |n\rangle$$

⇒ n-th excited state

$$E_n = (n + \frac{1}{2}) \hbar \omega$$



$$\left[\frac{(\tilde{a}^\dagger)^n}{\sqrt{n!}} \right] |0\rangle \equiv |n\rangle$$

$$n=0, 1, 2, \dots$$

* matrix representation in the basis of $\{|n\rangle\}$.

$$\langle n' | \tilde{a} | n \rangle = \sqrt{n} \delta_{n', n-1}, \quad \langle n' | \tilde{a}^\dagger | n \rangle = \sqrt{n+1} \delta_{n', n+1}$$


• \tilde{x} and \tilde{p} ?

$$\tilde{x} = \sqrt{\frac{\hbar}{2m\omega}} (\tilde{a} + \tilde{a}^\dagger)$$

$$\tilde{p} = i \sqrt{\frac{m\hbar\omega}{2}} (-\tilde{a} + \tilde{a}^\dagger)$$

$$\Rightarrow \langle n' | \tilde{x} | n \rangle = \sqrt{\frac{\hbar}{2m\omega}} (\sqrt{n} \delta_{n', n-1} + \sqrt{n+1} \delta_{n', n+1})$$

$$\langle n' | \tilde{p} | n \rangle = i \sqrt{\frac{m\hbar\omega}{2}} (-\sqrt{n} \delta_{n', n-1} + \sqrt{n+1} \delta_{n', n+1})$$



- Energy eigenfunction in x -space.

- ground-state wave function $\langle x|0\rangle$:

Ground-state ket $|0\rangle$ satisfies

$$\tilde{a}|0\rangle = 0 \Rightarrow \langle x|\tilde{a}|0\rangle = \sqrt{\frac{m\omega}{2\hbar}} \langle x|\tilde{x} + i\frac{\tilde{p}}{m\omega}|0\rangle$$

$$\Rightarrow \left(x + x_0^2 \frac{\partial}{\partial x} \right) \langle x|0\rangle = 0 \quad \left\| \quad x_0^2 = \frac{\hbar}{m\omega} \right.$$

1st. order diff. eq.

$$\langle x|0\rangle = A e^{-\frac{1}{2} \frac{x^2}{x_0^2}} \oplus \text{normalization } \int_{-\infty}^{\infty} dx \langle x|0\rangle = 1$$

$$= \frac{1}{\pi^{1/4} \sqrt{x_0}} \exp \left[-\frac{1}{2} \left(\frac{x}{x_0} \right)^2 \right]$$

- excited states.

$$\begin{aligned} \langle x|1\rangle &= \langle x|\tilde{a}^+|0\rangle = \sqrt{\frac{m\omega}{2\hbar}} \langle x|\tilde{x} - i\frac{\tilde{p}}{m\omega}|0\rangle \\ &= \frac{1}{\sqrt{2}x_0} \left(x - x_0^2 \frac{\partial}{\partial x} \right) \langle x|0\rangle \\ &\vdots \end{aligned}$$

$$\begin{aligned} \langle x|n\rangle &= \frac{1}{\sqrt{n!}} \langle x|(\tilde{a}^+)^n|0\rangle = \frac{1}{\sqrt{n!}} \left(\frac{1}{\sqrt{2}x_0} \right)^n \left(x - x_0^2 \frac{\partial}{\partial x} \right)^n \langle x|0\rangle \\ &= \frac{1}{\pi^{1/4} \sqrt{2^n n!}} \left(\frac{1}{x_0^{n+\frac{1}{2}}} \right) \left(x - x_0^2 \frac{\partial}{\partial x} \right)^n \exp \left(-\frac{1}{2} \left(\frac{x}{x_0} \right)^2 \right) \end{aligned}$$

- Let's check up what we know from the uncertainty principle.


$$\Delta x \Delta p \geq \frac{\hbar}{2}$$

- classical Hamiltonian :

$$H = \frac{p^2}{2m} + \frac{1}{2} m \omega^2 x^2 = E \quad (\text{Energy})$$

We know $\Delta p = p$, $\Delta x = x$ since

it's an oscillator.



$$E = \frac{\Delta p^2}{2m} + \frac{1}{2} m \omega^2 \Delta x^2 \geq \frac{\hbar^2}{8m} \Delta x^{-2} + \frac{1}{2} m \omega^2 \Delta x^2$$

has minimum at $\Delta x^2 = \frac{\hbar}{2m\omega}$

$$(E_{\min} = \frac{\hbar}{2} \omega)$$

$$\geq \frac{1}{4} \hbar \omega + \frac{1}{4} \hbar \omega$$

$$\Delta p^2 = \frac{\hbar m \omega}{2}$$

$$\geq \frac{1}{2} \hbar \omega \equiv E_{\text{ground}} \quad \circ$$

- Let's verify these steps with $\langle n | \circ \rangle$.

$$\langle \tilde{x} \rangle = 0, \quad \langle \tilde{p} \rangle = 0.$$

$$\begin{aligned} \langle \tilde{x}^2 \rangle &= \left\langle \frac{\hbar}{2m\omega} (\tilde{a} + \tilde{a}^\dagger)^2 \right\rangle = \frac{\hbar}{2m\omega} \langle 0 | (\tilde{a}^2 + \tilde{a}^{\dagger 2} + \tilde{a}^\dagger \tilde{a} + \tilde{a} \tilde{a}^\dagger) | 0 \rangle \\ &= \frac{\hbar}{2m\omega} \end{aligned}$$

\downarrow
 $1 + \tilde{a}^\dagger \tilde{a}$

$$\text{Similarly, } \langle \tilde{p}^2 \rangle = \frac{\hbar m \omega}{2} \quad \Leftrightarrow \quad \begin{cases} \Delta x^2 = \frac{\hbar}{2m\omega} \\ \Delta p^2 = \frac{\hbar m \omega}{2} \end{cases}$$

$$\left\langle \frac{\tilde{p}^2}{2m} \right\rangle = \frac{1}{4} \hbar \omega, \quad \left\langle \frac{1}{2} m \omega^2 \tilde{x}^2 \right\rangle = \frac{1}{4} \hbar \omega$$

$$E = \langle H \rangle = \frac{1}{2} \hbar \omega.$$

$$\langle (\Delta \tilde{x})^2 \rangle \langle (\Delta \tilde{p})^2 \rangle = \frac{\hbar}{2m\omega} \cdot \frac{\hbar m \omega}{2} = \frac{\hbar^2}{4}$$

for the n th state,

$$\langle (\Delta \tilde{x})^2 \rangle \langle (\Delta \tilde{p})^2 \rangle = \left(n + \frac{1}{2}\right)^2 \hbar^2 \geq \frac{\hbar^2}{4}$$

(2) Time Development of the Oscillation

24

- Heisenberg picture

$$\text{EOM: } \frac{d\tilde{p}(t)}{dt} = \frac{1}{i\hbar} [\tilde{p}(t), H]$$

$$\begin{cases} p(t) \equiv p^{(H)}(t) \\ x(t) \equiv x^{(H)}(t) \end{cases}$$

$$\frac{d\tilde{x}(t)}{dt} = \frac{1}{i\hbar} [\tilde{x}(t), H]$$

NOTE: We know that $\langle H \rangle$ is conserved, thus, $H(t) = H$.
But, let's just try with $H(t)$. (t-indep.)

$$H = \frac{\tilde{p}(t)^2}{2m} + \frac{1}{2}m\omega^2 \tilde{x}(t)^2$$

$$\Rightarrow \begin{cases} \frac{d\tilde{p}(t)}{dt} = -m\omega^2 \tilde{x}(t) \\ \frac{d\tilde{x}(t)}{dt} = \frac{p(t)}{m} \end{cases}$$

* Show $[x(t), p(t)] = i\hbar$
when $x(t) = e^{i\frac{H}{\hbar}t} x e^{-i\frac{H}{\hbar}t}$
 $p(t) = e^{i\frac{H}{\hbar}t} p e^{-i\frac{H}{\hbar}t}$

$$\Rightarrow \frac{d}{dt} \begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix} = \begin{pmatrix} 0 & \frac{1}{m} \\ -m\omega^2 & 0 \end{pmatrix} \begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix}$$

diagonalization:
 $\equiv A$

eigenvalues

$$\begin{cases} i\omega \leftarrow \begin{pmatrix} -\frac{i}{m\omega} \\ 1 \end{pmatrix} \\ -i\omega \leftarrow \begin{pmatrix} 1 \\ -im\omega \end{pmatrix} \end{cases}$$

\hookrightarrow

$$X^{-1} A X = D$$

$$D = \begin{pmatrix} i\omega & 0 \\ 0 & -i\omega \end{pmatrix}$$

\hookrightarrow

$$A = X D X^{-1}$$

$$X = \frac{1}{\sqrt{2}} \begin{pmatrix} -\frac{i}{m\omega} & 1 \\ 1 & -im\omega \end{pmatrix}$$

$$\Rightarrow \frac{d}{dt} \begin{bmatrix} X^{-1} \begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix} \end{bmatrix} = D \begin{bmatrix} X^{-1} \begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix} \end{bmatrix}$$

$$X^{-1} \begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \hat{m}\omega & 1 \\ 1 & \frac{\hat{p}}{m\omega} \end{pmatrix} \begin{pmatrix} \tilde{x}(t) \\ \tilde{p}(t) \end{pmatrix}$$

$$= \frac{1}{\sqrt{2}} \begin{pmatrix} \hat{m}\omega \tilde{x}(t) + \tilde{p}(t) \\ \tilde{x}(t) + \frac{\hat{p}}{m\omega} \tilde{p}(t) \end{pmatrix} \equiv A \begin{pmatrix} \hat{\psi}_1(t) \\ \hat{\psi}_2(t) \end{pmatrix}$$

$$\Rightarrow \frac{d}{dt} \begin{pmatrix} \hat{\psi}_1(t) \\ \hat{\psi}_2(t) \end{pmatrix} = \begin{pmatrix} \hat{i}\omega & 0 \\ 0 & -\hat{i}\omega \end{pmatrix} \begin{pmatrix} \hat{\psi}_1(t) \\ \hat{\psi}_2(t) \end{pmatrix}$$

Any diagonal matrix with c-numbers,

Choose it to be "I".

$$\therefore \begin{cases} \hat{\psi}_1(t) = c_1 e^{i\omega t} \\ \hat{\psi}_2(t) = c_2 e^{-i\omega t} \end{cases} \quad \parallel \quad \begin{cases} c_1 = \hat{\psi}_1(0) \\ c_2 = \hat{\psi}_2(0) \end{cases}$$

also for $\tilde{a}^\dagger, \tilde{a}$

time-invariant : $\psi_1(t) \cdot \psi_2(t) = c_1 c_2$.

$$\Rightarrow \left[\tilde{x}(t) - \frac{\hat{p}}{m\omega} \tilde{p}(t) \right] = \left[\tilde{x}(0) - \frac{\hat{p}}{m\omega} \tilde{p}(0) \right] e^{i\omega t}$$

$$\left[\tilde{x}(t) + \frac{\hat{p}}{m\omega} \tilde{p}(t) \right] = \left[\tilde{x}(0) + \frac{\hat{p}}{m\omega} \tilde{p}(0) \right] e^{-i\omega t}$$

Q. Is it really classical?

$$\Rightarrow \begin{cases} \tilde{x}(t) = \tilde{x}(0) \cos \omega t + \frac{\tilde{p}(0)}{m\omega} \sin \omega t \\ \tilde{p}(t) = -m\omega \tilde{x}(0) \sin \omega t + \tilde{p}(0) \cos \omega t \end{cases}$$

$$\begin{cases} \cos \omega t = \frac{e^{i\omega t} + e^{-i\omega t}}{2} \\ \sin \omega t = \frac{e^{i\omega t} - e^{-i\omega t}}{2i} \end{cases}$$

* What about $\tilde{a}, \tilde{a}^\dagger$?

Note that, $\hat{\psi}_1 = \frac{1}{\sqrt{2}} (\hat{m}\omega \tilde{x} + \tilde{p})$

$$\propto \tilde{a}^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(\tilde{x} - \frac{i\tilde{p}}{m\omega} \right)$$

and $\hat{\psi}_2 \propto \tilde{a}$; $\Rightarrow \begin{cases} \tilde{a}^\dagger(t) = e^{i\omega t} \tilde{a}^\dagger(0) \\ \tilde{a}(t) = e^{-i\omega t} \tilde{a}(0) \end{cases}$

These can be verified by using

the Baker-Hausdorff

Lemma : $A(t) = e^{\frac{i\hbar t}{2}} A e^{-\frac{i\hbar t}{2}}$

= ... (pp 95.)
of Sakurai.

$\therefore \tilde{a}^\dagger(t) \tilde{a}(t) = \text{time-invariant} ; [H, \tilde{a}^\dagger \tilde{a}] = 0$

\rightarrow simultaneous eigenket !!!

Look at :

$$\begin{cases} \tilde{x}(t) = \tilde{x}(0) \cos \omega t + \frac{\tilde{p}(0)}{m\omega} \sin \omega t \\ \tilde{p}(t) = -m\omega \tilde{x}(0) \sin \omega t + \tilde{p}(0) \cos \omega t \end{cases}$$

$$\Leftrightarrow m \frac{d\tilde{x}}{dt} = \tilde{p}, \text{ just like C.M.}$$

But, $\langle \tilde{x} \rangle$ and $\langle \tilde{p} \rangle$ are "not" oscillating. !!!

although $\tilde{x}(t)$ and $\tilde{p}(t)$ look like oscillating.

|| NOTE: $\langle \tilde{x} \rangle = 0$, $\langle \tilde{p} \rangle = 0$, for all $|n\rangle$.

Q. Can we find a "Quantum" state that behaves just like classical $\langle x \rangle$ and $\langle p \rangle$?

* Coherent States \leftarrow This is the one.

Why do we need this?

- We live in a "classical" world,
But we want to control a "Quantum" world.
: We need a "bridge"!

the easiest way to make the coherent state



$$|S_0\rangle = J(S_0) |0\rangle$$

move the ground state to S_0 .

wave function $\psi_{S_0}(x) = \psi_0(x - S_0)$ $\parallel \begin{cases} \langle x | J(S_0) \\ = \langle x - S_0 | \end{cases}$

observables :

$$\langle S_0 | \tilde{x} | S_0 \rangle = \langle 0 | J^\dagger(S_0) \tilde{x} J(S_0) | 0 \rangle = \underline{S_0}$$

$$\langle S_0 | \tilde{p} | S_0 \rangle = 0$$

$$\langle S_0 | H | S_0 \rangle = \langle 0 | \frac{\tilde{p}^2}{2m} | 0 \rangle + \frac{1}{2} m \omega^2 \langle 0 | (\tilde{x} + S_0)^2 | 0 \rangle$$